arXiv:1003.3804v2 [math.AG] 23 Jun 2011

ON THE CANONICAL DEGREES OF CURVES IN VARIETIES OF GENERAL TYPE

by

Pascal Autissier, Antoine Chambert-Loir & Carlo Gasbarri

Introduction

In this article, we work over a fixed algebraically closed field k of characteristic zero. The word *variety* means an integral k-scheme of finite type, and a *curve* is a variety of dimension 1. If C is a projective curve, we let g_C be its geometric genus and $\chi(C) = 2 - 2g_C$ its geometric Euler characteristic; we also let $\deg_C L$ be the degree of a line bundle L on C.

A widely believed conjecture predicts that curves of bounded geometric genus lying on a variety of general type X form a bounded family. An effective version of this conjecture can be stated in the following way:

Conjecture 1. — *Let X be a smooth projective variety of general type; let K_X be the canonical line bundle of X. Then there exist real numbers A and B, and a strict closed subscheme $Z \subset X$ such that for every projective curve C and any finite morphism $g\colon C \to X$ such that $g(C) \not\subset Z$, we have*

$$\deg_C(g^*K_X) \leq A(-\chi(C)) + B\deg(g).$$

When $\dim(X) = 1$ the conjecture is true, and is a straightforward consequence of the Riemann–Hurwitz formula. Concerning surfaces, the most important result in its direction is a now well-known theorem by Bogomolov [**3**] (and its further developments) who proves such a statement under the additional assumption that X is a minimal surface with positive Segre class. Over the complex numbers, and under the supplementary assumption that X is Brody-hyperbolic, *i.e.,* that any holomorphic map from $\mathbf{C}$ to X is constant, Demailly established this conjecture in [**7**] (Theorem 2.1); he shows that one may take $Z = \emptyset$ and $B = 0$.

Of course, an effective geometric upper-bound of the real numbers A and B of the conjecture would be of the utmost interest. Besides the very interesting paper [**11**] of Miyaoka, no method is known to understand the constant B. On the other side, many interesting conjectures and guesses have been made concerning the constant A and its meaning. For example, if X is a (smooth projective) minimal surface of general type, that article [**11**] explicits, for any positive real number ε, a real number B such that every *smooth* projective curve C contained in X satisfies

$$\deg_C K_X \leq (\frac{3}{2}+\varepsilon)(-\chi(C))+B.$$

Exploiting the analogy between number theory, Nevanlinna theory and algebraic geometry, P. Vojta [**18**] proposed very deep and ambitious conjectures bounding the canonical height of algebraic points of bounded degree in fibrations. The first version of his conjectures suggests that for any positive real number ε, conjecture 1 could hold with $A = \dim(X) + \varepsilon$, the constant B and the exceptional locus Z depending on ε, provided one restricts oneself to curves of given gonality. (Recall that the gonality of a curve is the least degree of a non-constant morphism to the projective line; obviously, horizontal curves of bounded degree in a fibration have bounded gonality.) A more recent version of his conjecture, published in [**19**] replaces $A = \dim X + \varepsilon$ by the more optimistic $A = 1 + \varepsilon$, again with the same restriction on gonality.

More prudently, McQuillan [**9**] asked whether the constant A could be related to the holomorphic sectional curvature of the variety X. In this direction, we prove the following theorem, whose proof relies indeed on the computation by Tashiro of the holomorphic sectional curvatures of bounded symmetric domains ([**15, 16**]), see Lemma 7.

Theorem 2. — *Let M be a bounded hermitian domain whose irreducible factors are of classical type and let X be a compact quotient of M. Then, for any projective curve C and any finite morphism $g\colon C \to X$,*

$$\deg_C(g^*K_X) \leq -\dim(X)\chi(C).$$

In other words, we establish Conjecture 1 for such varieties, with $A = \dim(X)$.

Conversely, and this was the original motivation of this note, we also want to indicate, by very elementary examples, that one cannot go under $A = \dim(X)$ in general, or, in other words, that Vojta's refined expectation cannot hold without the above-mentioned restriction on the gonality. Namely, we shall prove the following theorem.

Theorem 3. — *Let d be an integer such that $d \geq 2$. There exists a smooth projective variety X of dimension d whose canonical bundle K_X is ample and* *a family* (C_n, f_n)*, where* C_n *is a smooth projective curve and* $f_n\colon C_n \to X$ *is a finite morphism satisfying the following properties:*

a) $\inf_n \dfrac{\deg_{C_n} f_n^* K_X}{-\chi(C_n)} \geq d$*;*
b) $\lim_n(-\chi(C_n)) = +\infty$*;*
c) *the union of their images* $f_n(C_n)$ *is dense in* X *for the Zariski topology.*

A slightly weaker form of Theorem 3 will be consequence of well known facts on modular curves and Hecke correspondences: in our example, X is a power of a modular curve $X_0(p)$, for a large prime p, and the curves C_n are modular curves $X_0(p\ell_1 \dots \ell_{d-1})$, defined by distinct prime numbers $\ell_1, \dots, \ell_{d-1}$. However, this example only establishes a slight weakening of Theorem 3, where d is replaced by $d - \varepsilon$, for arbitrary positive real number ε.

As we realized later on, considering *compact* Shimura curves instead of standard modular curves allows to prove Theorem 3 as it stands.

Note also that this theorem does not contradict Miyaoka's result [**11**] mentioned above, because there is no reason that the curves $f_n(C_n)$ in X are smooth. However, in our examples, the morphism f_n will be an immersion in the sense of differential geometry, *i.e.*, its tangent map is injective.

We view these examples as a nice implementation of McQuillan's principle that *Shimura varieties are a rich source of examples for Diophantine geometry.* Observe that they do not contradict Vojta's expectations since he restricts himself to curves in a fibration which have bounded degree. The gonality of such curves is obviously bounded, while, according to a theorem of Abramovich [**1**], the gonality of modular curves tends to infinity.

In fact, using weak approximation, we show (Theorem 8) that one may take for X any compact Shimura variety associated to any **Q**-anisotropic, semisimple, simply connected **Q**-group **G**, provided X contains at least one totally geodesic compact Riemann surface.

Acknowledgments: This work began during an instructional conference on Diophantine Geometry, organized by two of us in University of Rennes under the auspices of Centre Émile Borel and Institut Henri Poincaré. We thank these organizations to have given us the opportunity to discuss this subject.

We also thank B. Edixhoven, A. Levin, M. McQuillan, E. Ullmo and P. Vojta for their interest and their comments.

The three authors are supported by the project *Positive* of Agence nationale de la recherche, under agreement ANR-2010-BLAN-0119-01.

1. Modular curves: proof of Theorem 3$-\varepsilon$

Let N be a positive integer. The (affine) modular curve $Y_0(N)$ classifies couples (E,G), where E is an elliptic curve and $G\subset E$ is a cyclic subgroup of order N. Precisely, its k-points are in natural bijection with the set of isomorphism classes of such couples (E,G), where E is an elliptic curve over k. The modular curve $X_0(N)$ is the unique smooth projective curve containing $Y_0(N)$ as a dense open subset.

Let N be a positive integer and let M be a divisor of N such that $M<N$. Let (E,G) be a pair consisting of an elliptic curve over k and a cyclic subgroup of order N. Then $G_M=G\cap E[M]$ is a cyclic subgroup of E of order M and the pair (E,G_M) defines a k-point of $Y_0(M)$. We let s_M^N be the unique morphism of $X_0(N)$ to $X_0(M)$ such that $s_M^N(E,G)=(E,G_M)$. The maps s_M^N satisfy the transitivity property $s_M^N=s_M^{M'}\circ s_{M'}^N$ whenever M,M' and N are positive integers such that $M|M'|N$. Recall that for any elliptic curve E and any positive integer N, there exist exactly $N\prod_{\ell|N}(1+\ell^{-1})$ cyclic subgroups of order N, where the product ranges over the prime numbers dividing N. Consequently, the degree of s_1^N is equal to $N\prod_{\ell|N}(1+\ell^{-1})$. In particular, when M and N/M are coprime, the degree of s_M^N is equal to $(N/M)\prod_{\ell|N/M}(1+\ell^{-1})$.

On the other hand, the quotient curve $E/G_{N/M}$ together with the image $G/G_{N/M}$ of G again defines a point of $Y_0(M)$. We let $t_M^N\colon X_0(N)\to X_0(M)$ be the unique morphism of curves such that $t_M^N(E,G)=(E/G_{N/M},G/G_{N/M})$. The same argument as before, now applied to the isogeny dual to the projection $E\to E/G_{N/M}$, implies that the degree of t_M^N is equal to that of s_M^N.

From now on, we assume that N is the product of r distinct prime numbers. The formulae for the degrees of s_M^N and t_M^N become

$$\deg(s_M^N)=\prod_{\ell|N/M}(\ell+1). \tag{1}$$

Moreover, according to [**14**], Prop. 1.40 p. 23 and Prop. 1.43 p. 24, the Euler characteristic of $X_0(N)$ satisfies the inequality

$$0\leq\frac{1}{6}\prod_{\ell|N}(\ell+1)+\chi(X_0(N))\leq\frac{13}{6}2^r. \tag{2}$$

Moreover, $\chi(X_0(N))<0$ for $N\geq 22$.

Let $p,\ell_1,\dots,\ell_{d-1}$ be distinct prime numbers, with $p\geq 23$. Let us pose $N=p\ell_1\dots\ell_{d-1}$. Let X be the variety $X_0(p)^d$, let C_N be the curve $X_0(\ell_1\dots\ell_{d-1}p)$ and let $f_N\colon C_N\to X$ be the morphism given by $f_N=(g_1,\dots,g_{d-1},g_d)$, where $g_d=s_p^N$ and $g_i=t_p^{\ell_i p}\circ s_{\ell_i p}^N$ for $1\leq i\leq d-1$.

By Equation (2), the Euler characteristics of the curves C_N tend to infinity. Therefore, choosing arbitrary large prime numbers p, and up to arbitrary small ε, Theorem 3 is a direct consequence of the two lemmas below.

Lemma 4. — *The curve (C_N, f_N) satisfies the inequality*

$$\deg f_N^* K_X \geq d\left(1 - \frac{26}{p+1}\right)(-\chi(C_N)).$$

Proof. — For $1 \leq i \leq d$, let $\mathrm{pr}_i \colon X \to X_0(p)$ be the ith projection; one has $K_X = \sum_{i=1}^d \mathrm{pr}_i^* K_{X_0(p)}$ hence

$$\deg f_N^* K_X = \sum_{i=1}^{d} \deg g_i^* K_{X_0(p)} = -\sum_{i=1}^{d} \deg(g_i)\chi(X_0(p)).$$

By Equation (1),

$$\deg(g_i) = (\ell_1 + 1)\dots(\ell_{d-1}+1),$$

hence

$$\deg(f_N^* K_X) = -d(\ell_1+1)\dots(\ell_{d-1}+1)\chi(X_0(p)).$$

Since Equation (2) implies that

$$-\chi(X_0(p)) \geq \frac{p+1}{6} - \frac{13}{3},$$

we obtain the inequality

$$\deg(f_N^* K_X) \geq d(p-25)(\ell_1+1)\dots(\ell_{d-1}+1).$$

On the other hand, it again follows from Equation (2) that

$$-\chi(C_N) \leq \frac{1}{6}(p+1)(\ell_1+1)\dots(\ell_{d-1}+1).$$

Consequently,

$$\frac{\deg(f_N^* K_X)}{-\chi(C_N)} \geq d\frac{p-25}{p+1} = d\left(1 - \frac{26}{p+1}\right),$$

as was claimed. ☐

Lemma 5. — *Let p and d be fixed. Then the union of the curves $f_N(C_N)$ is dense for the Zariski topology, where N varies among the set of of the form $N = \ell_1 \dots \ell_{d-1} p$ as above.*

Proof. — The proof is by induction on d. When $d = 1$, it clearly holds since $C_p = X_0(p)$ and $f_p \colon X_0(p) \to X$ is the identity map. Let us assume that the result holds for $d-1$ and let us prove it for d. Assume by contradiction that there exists a divisor H in X containing the images of all curves C_N above.

Let $X' = X_0(p)^{d-1}$ and let $q \colon X \to X'$, $(x_1, \dots, x_d) \mapsto (x_2, \dots, x_d)$, be the projection obtained by forgetting the first component. For any integer N'

of the form $\ell_2 \dots \ell_{d-1} p$ which is the product of distinct prime numbers, let $C_{N'} = X_0(N')$ and $f'_{N'} \colon C_{N'} \to X'$ be the corresponding morphism. If ℓ_1 is any prime number not dividing N', there is a commutative diagram

$$\begin{tikzcd}
X_0(N) \arrow[r, "f_N"] \arrow[d, "s^N_{N'}"] & X \arrow[d, "q"] \\
X_0(N') \arrow[r, "f'_{N'}"] & X',
\end{tikzcd}$$

where $N = \ell_1 N'$. By induction, all curves $f'_{N'}(X_0(N'))$ are contained in $q(H)$, so that $q(H) = X'$.

The k-points in $X_0(N')$ of the form (E, G) where E is an elliptic curve without complex multiplication are Zariski dense; consequently, the same holds for their images in X' by the morphisms $f'_{N'}$. Since $\dim H = \dim X'$ and the restriction q_H of q to H is surjective, there exists such a point $f'_{N'}(E, G)$ above which q_H is finite.

For any prime number ℓ which does not divide N', let K_ℓ be a cyclic subgroup of order ℓ in E. The subgroup $K_\ell + G$ of E is cyclic of order $N = \ell N'$ and the point $f_N(E, K_\ell + G)$ of X is a point of H mapping to $f'_{N'}(E, G)$ by q. Moreover, $\mathrm{pr}_1(f_N(E, K_\ell + G)) = g_1(E, K_\ell + G) = (E/K_\ell, (K_\ell + G_p)/K_\ell)$. However, if ℓ and ℓ' are distinct prime numbers not dividing N', the elliptic curves E/K_ℓ and $E/K_{\ell'}$ are not isomorphic, since E has no complex multiplication (otherwise, E would admit an isogeny of degree $\ell\ell'$). Consequently, the fibre of q_H above $f'_{N'}(E, G)$ is infinite, contradiction. □

2. Products of compact Shimura curves

In this section, we briefly discuss on how the consideration of compact Shimura curve allows to prove Theorem 3 as we stated it. Our main references are [**10, 5**]; see also [**17, 4**].

Let D be an indefinite quaternion algebra over $\mathbf{Q}$ of reduced discriminant $\delta > 1$, viewed as a subalgebra of $\mathrm{M}_2(\mathbf{R})$; let R be a fixed maximal order in D (they are all conjugate) and let Γ_R be the subgroup $R^* \cap \mathrm{SL}_2(\mathbf{R})$ of $\mathrm{GL}_2(\mathbf{R})$. Elements of Γ_R act by homographies on the Poincaré upper-half-plane $\mathfrak{h}$. The Shimura curve C associated to D is the quotient of $\mathfrak{h}$ by Γ_R. It is a compact algebraic complex curve whose points are in one-to-one correspondence with the set of (isomorphism classes of) complex abelian surfaces endowed with an action of R, also called *false elliptic curves.* These abelian surfaces carry a canonical principal polarization once an element of square $-\delta$ is fixed in R.

See also [**5**, p. 594] for an adelic description of this Shimura curve, in the spirit of the general theory of Shimura varieties, as well as for its coverings induced by level-structures on these abelian surfaces. For any square free positive integer N which is prime to δ, one has in particular a curve C_N parametrizing isogenies of *false degree* N between *false elliptic curves*, that is isogenies $f\colon A \to A'$ commuting with the action of R such that the composite $f^\vee \circ f$ of f with its dual isogeny $f^\vee$ is the multiplication by N. There are natural maps $s^N_M\colon C_N \to C_M$ and $t^N_M\colon C_N \to C_M$ between these curves, whenever M divides N. If $N \geq 4$, this curve is a *fine* moduli space; equivalently, $C_N(\mathbf{C})$ is the quotient of the upper-half-plane by a discrete, cocompact, and torsion free subgroup of $\mathrm{SL}_2(\mathbf{R})$. Since there are no cusps at infinity, the complex description of these curves, or their moduli interpretation, shows that these maps are finite and *étale*.

We may now replace the modular curves of Section 1 by these compact Shimura curves. Our claim is that the curves (C_N, f_N) we obtain satisfy the statement of Theorem 3. Here, N is the product of d distinct prime numbers not dividing δ, among which a fixed prime number $p > 4$, and f_N is a morphism from C_N to $X = C_p^d$. Since there are no points at infinity, Lemma 4 becomes

Lemma 6. — *The curve* (C_N, f_N) *satisfies the equality*

$$\deg f_N^* K_X = -d\chi(C_N).$$

Lemma 5 continues to hold. Since we made use of elliptic curves without complex multiplication, its proof needs to be adapted. However, it suffices to observe that given any prime number ℓ which does not divide δ, there are infinitely many prime numbers ℓ' such that our false elliptic curves have no isogeny of false degree $\ell\ell'$. (Otherwise, almost all prime numbers would be reduced norms in D.)

3. Compact quotients of hermitian symmetric domains

3.1. Hermitian symmetric spaces; Bergman metric. — Let M be a Hermitian symmetric space, namely a connected Kähler complex manifold such that each point of M is an isolated fixed point of an involutive holomorphic isometry of M. Set $m = \dim(M)$.

Following the notation of [**8**] (chap. VIII, §4), let $A(M)$ be the group of holomorphic isometries of M and let $G = A_0(M)$ be its identity component; this is a real connected Lie group. We fix a point $o \in M$ and consider its stabilizer K in $A_0(M)$; this is a compact subgroup; as a real manifold, M can be identified to the homogeneous space G/K.

We shall always assume that M is of noncompact type: G is semisimple and K is a maximal compact subgroup of G; then M is simply connected ([**8**], p. 376, Theorem 4.6) and there exists ([**8**], p. 382, Theorem 7.1) a holomorphic diffeomorphism from M to a bounded symmetric domain D_M in the complex space $\mathbf{C}^m$, "symmetric" meaning that each point of D_M is an isolated fixed point of an involutive holomorphic diffeomorphism of D_M onto itself.

The Bergman kernel of D_M is the function on $D_M \times D_M$ given by

$$K(z,w)=\sum_{n=0}^{\infty}\varphi_n(z)\overline{\varphi_n(w)},$$

where (φ_n) is any orthonormal basis of the Hilbert space $\mathscr{H}(D_M)\cap \mathrm{L}^2(D_M)$ (the convergence is uniform on each compact subset of $D_M \times D_M$). The Bergman metric is the Hermitian metric on the tangent bundle $\mathrm{T}D_M$ defined as

$$H=2\sum_{1\leq i,j\leq m}\frac{\partial}{\partial z_i}\frac{\partial}{\partial \bar z_j}\log K(z,z)\mathrm{d}z_i\otimes \mathrm{d}\bar z_j.$$

Any holomorphic diffeomorphism of D_M is an isometry with respect to that metric ([**8**], p. 370, Prop. 3.5). Moreover, this is a Kähler metric, with associate Kähler form

$$\omega_M = i\partial\bar\partial \log K(z,z).$$

We assume, as we may, that the Kähler structure of M is the pull-back of the Bergman metric on D_M by some (hence, any) holomorphic diffeomorphism of M to D_M. It induces a metric $\|\cdot\|_{\mathrm{B}}$ on the canonical bundle $K_M=\bigwedge^m\Omega^1_M$ of M. Proposition 3.6 (p. 371) of [**8**] implies the following relation between the Kähler form of M and the first Chern form on K_M:

$$\omega = 2\pi c_1(K_M,\|\cdot\|_{\mathrm{B}}). \tag{3}$$

In other words, the Bergman metric on D_M is Kähler-Einstein.

3.2. Holomorphic sectional curvature. — Let us recall a definition of the holomorphic sectional curvature. Let $p\in M$ and let us consider holomorphic maps $\gamma\colon U\to M$, where U is the open unit disk in $\mathbf{C}$, such that $\gamma(0)=0$ and $\gamma'(0)\neq 0$. Let us write $\gamma^*\omega = F_\gamma(z)i\mathrm{d}z\wedge\mathrm{d}\bar z$ on U, where F_γ is nonnegative, $F_\gamma(0)\neq 0$. In a neighborhood of 0, the form $\gamma^*\omega$ defines a metric on the canonical bundle K_U and its curvature form is given by

$$R_\gamma=\frac{i}{2\pi}\partial\bar\partial\log(F_\gamma).$$

The Ricci curvature Ric_γ of $\gamma^*\omega$ is defined by the formula

$$\mathrm{Ric}_\gamma(z)F_\gamma(z)i\mathrm{d}z\wedge\mathrm{d}\bar z=-i\partial\bar\partial\log(F_\gamma).$$

This is a function on a neighborhood of 0 in U, and its value at $z=0$ only depends on the tangent line at p generated by $\gamma'(0)$.

The holomorphic sectional curvature $S_M(p)$ at p is then defined as

$$S_M(p)=\sup_{\substack{\gamma\colon U\to M\\ \gamma(0)=p}}\mathrm{Ric}_\gamma(0) \tag{4}$$

Since the group of holomorphic isometries of M acts transitively on M, the holomorphic sectional curvature is actually independent of p; we shall denote it by S_M. Since M is a Hermitian symmetric space of the non-compact type ([**8**], Ch. VIII, Theorem 7.1, p. 382), we know that S_M is negative ([**8**], Ch. V, Theorem 3.1, p. 241).

***Lemma 7* (Tashiro).** — *Assume that the irreducible factors of M are of classical type. Then*

$$\frac{-1}{S_M} \leq \dim(M).$$

Proof. — The proof follows from the explicit computations in [**15, 16**].

If M_1 and M_2 are complex hermitian manifolds the holomorphic sectional curvatures of which satisfy $-1/S_1 \leq \dim(M_1)$ and $-1/S_2 \leq \dim(M_2)$, then the holomorphic sectional curvature of $M = M_1 \times M_2$ is bounded from above by

$$S_M \leq -\frac{\dim(M_1)^{-1}\dim(M_2)^{-1}}{\dim(M_1)^{-1}+\dim(M_2)^{-1}} = -\frac{1}{\dim(M_1)+\dim(M_2)}.$$

Consequently, it suffices to treat the case of an irreducible domain belonging to one of the four classical types.

Type I(n,m). — *M is the set of matrices Z of type (n,m), where $n \geq m \geq 1$, such that* $\mathrm{I}_m - Z^*Z$ *is positive definite.* Formula (6.3) of [**16**] shows that

$$\frac{-1}{S_M} = \sup_{Z\in \mathrm{Mat_{n,m}}(\mathbf{C})} \frac{n+m}{2}\frac{(\mathrm{Tr}(Z^*Z))^2}{\mathrm{Tr}(Z^*ZZ^*Z)}.$$

The matrix $H = Z^*Z$ is hermitian of size m; let $\lambda_1, \dots, \lambda_m$ be its eigenvalues. Applying the Cauchy-Schwarz inequality, we obtain

$$(\mathrm{Tr}(Z^*Z))^2 = \left(\sum_{j=1}^m \lambda_j\right)^2 \leq m\sum_{j=1}^m \lambda_j^2 = m\,\mathrm{Tr}(Z^*ZZ^*Z); \tag{5}$$

equality holds if Z^*Z is a homothety. Therefore,

$$\frac{-1}{S_M} = m(m+n)/2 \leq mn = \dim(M).$$

Type II(n). — *M is the set of symmetric matrices Z of size n such that* $\mathrm{I}_n - Z^*Z$ *is positive definite.* Formula (6.4) of [**16**] claims that

$$\frac{-1}{S_M} = \sup_{Z\in \mathrm{Sym}_n(\mathbf{C})} \frac{n+1}{2}\frac{(\mathrm{Tr}(Z^*Z))^2}{\mathrm{Tr}(Z^*ZZ^*Z)}.$$

It follows that

$$\frac{-1}{S_M} \leq n(n+1)/2 = \dim(M).$$

Type III(n). — *M is the set of skew-symmetric matrices Z of size n such that $\mathrm{I}_n - Z^*Z$ is positive definite.* Formula (6.5) of [**16**] claims that

$$\frac{-1}{S_M} = \sup_{Z\in\mathrm{Sym}_n(\mathbf{C})} \frac{n-1}{2}\frac{(\mathrm{Tr}(Z^*Z))^2}{\mathrm{Tr}(Z^*ZZ^*Z)}.$$

It follows that

$$\frac{-1}{S_M} \leq n(n-1)/2 = \dim(M).$$

Type IV(n). — *M is the set of matrices $Z = (z_1, \dots, z_n)$ of type $(n, 1)$ such that $|\sum z_j^2| < 1$ and $1 + |\sum z_j^2|^2 - 2\sum|z_j|^2 > 0$.* One has (see [**15**], page 75),

$$-S_M = \inf_{Z\in\mathbf{C}^n}\left(\frac{1}{n} + \frac{\sum|\bar{z}_j z_k - \bar{z}_k z_j|^2}{2n(\sum|z_j|^2)^2}\right) = \frac{1}{n}.$$

Again,

$$\frac{-1}{S_M} = n = \dim(M).$$

This concludes the proof. □

3.3. Compact quotients. — Let Γ be a cocompact discrete subgroup of G which acts without fixed points on M. Then, $X_\Gamma = \Gamma\backslash M$ is a compact connected complex manifold. The Kähler-Einstein structure descends to a Kähler-Einstein metric ω_Γ on X_Γ. The canonical line bundle K_{X_Γ} is naturally metrized; in view of Equation (3), its Chern form satisfies

$$\omega_\Gamma = 2\pi c_1(K_{X_\Gamma}, \|\cdot\|),$$

hence is positive. By Kodaira's theorem, X_Γ is a projective complex manifold.

3.4. Compact Riemann surfaces. — Let Y be a smooth compact Riemann surface and let $f\colon Y \to X_\Gamma$ be a finite morphism. One has

$$\int_Y f^*(\omega_\Gamma) = 2\pi \deg_Y(f^*K_{X_\Gamma}). \tag{6}$$

Moreover, the form $f^*(\omega_\Gamma)$ induces a hermitian metric on the canonical line bundle $K_Y(-\Delta_f)$ of Y, where Δ_f is the branching divisor of f. Let R_Y be its curvature, one has

$$\int_Y R_Y = \deg(K_Y - \Delta_f) = -\chi(Y) - \deg(\Delta_f).$$

By the definition of the holomorphic sectional curvature,

$$R_Y \geq \frac{-S_M}{2\pi} f^*(\omega_\Gamma).$$

This shows that

$$-\chi(Y) \geq -\chi(Y) - \deg(\Delta_f) \geq -S_M \deg_Y(f^*K_{X_\Gamma}). \tag{7}$$

Given Lemma 7, this establishes Theorem 2.

3.5. Totally geodesic Riemann surfaces. — Let (X,ω) be a Kähler complex manifold, let Y be a Riemann surface and let $f\colon Y\to X$ be a holomorphic immersion. We say that f is totally geodesic if any geodesic of X contained in Y is a geodesic of Y with respect to the Kähler form $f^*\omega$. This is equivalent to the vanishing of the second fundamental form of Y ([**8**], Theorem 14.5, p. 80) and implies that for each $p\in Y$, the Ricci curvature of $f^*\omega$ at p equals the holomorphic sectional curvature of X at $f(p)$. (This follows, for example, for Gauss equation relating the curvature tensor of a Riemannian manifold, that of a submanifold and the second fundamental form, see [**2**], Theorem 1.72.)

Let us assume that $X=X_\Gamma=\Gamma\backslash M$. Any immersion $f\colon Y\to X$ lifts to a holomorphic immersion $\tilde f\colon \mathbf{D}\to M$, where $\mathbf{D}$ is the open unit disk. The conditions for f and $\tilde f$ to be totally geodesic are equivalent. Moreover, since automorphisms of M preserve its Kähler form, we obtain that for any such automorphism g, $g\circ\tilde f$ is totally geodesic if and only if $\tilde f$ is.

For a totally geodesic map $f\colon Y\to X_\Gamma$, where Y is compact Riemann surface, inequality (7) becomes an equality:

$$-\chi(Y)=-S_M\deg_Y(f^*K_{X_\Gamma})\qquad (Y\text{ totally geodesic}). \tag{8}$$

3.6. Constructing one totally geodesic Riemann surface. — It does not seem easy to construct totally geodesic compact Riemann surfaces in arbitrary compact quotients of symmetric hermitian domains. However, let us show some examples, coming from the theory of Shimura varieties.

We restrict the situation to the case where the Lie group G is given by $G=\mathbf{G}(\mathbf{R})$, for some connected semisimple and simply connected $\mathbf{Q}$-*anisotropic* linear algebraic group $\mathbf{G}$ over $\mathbf{Q}$. Then, for any congruence subgroup Γ which acts without fixed point on $M=G/K$, the quotient space $X_\Gamma=\Gamma\backslash M$ is a smooth projective variety.

Let D_1 be a quaternion algebra over $\mathbf{Q}$ and let $\mathbf{G}_1=\mathrm{SL}(D_1)$ be the corresponding almost simple $\mathbf{Q}$-group. We assume that D_1 is not split, *i.e.*, is not isomorphic to $\mathrm{M}_2(\mathbf{Q})$ but that $D_1\otimes_{\mathbf{Q}}\mathbf{R}$ is split, so that $G_1=\mathbf{G}_1(\mathbf{R})\simeq\mathrm{SL}_2(\mathbf{R})$. Let $K_1\simeq\mathrm{SO}_2(\mathbf{R})$ be a maximal compact subgroup in G_1, let $M_1=G_1/K_1$; then M_1 is a hermitian symmetric domain, and is isomorphic to the Poincaré upper half-plane. Finally, let Γ_1 be a congruence subgroup in $G_1(\mathbf{Q})$, small enough so as to acts freely on M_1. Then $X_1=\Gamma_1\backslash M_1$ is a smooth projective curve.

Let F be a totally real field such that the simple F-algebra $D=D_1\otimes_{\mathbf{Q}}F$ is not split. Let $\mathbf{G}=\mathrm{SL}(D)$ be the corresponding almost simple F-group; by Weil's restriction of scalars, we view it as a semisimple $\mathbf{Q}$-group. One has $G=\mathbf{G}(\mathbf{R})\simeq\mathrm{SL}(2,\mathbf{R})^d$, where $d=[F:\mathbf{Q}]$. Similarly, let K be a maximal compact subgroup of G, let $M = G/K$ and let Γ be a congruence subgroup of $\mathbf{G}(\mathbf{Q})$ acting freely on M. Then, $X = \Gamma\backslash M$ is a smooth projective variety.

Moreover, the algebraic group $\mathbf{G}_1$ embeds naturally in $\mathbf{G}$; over $\mathbf{R}$, this is the diagonal embedding $\mathrm{SL}(2,\mathbf{R}) \hookrightarrow \mathrm{SL}(2,\mathbf{R})^d$. Let us assume that K contains K_1 and that Γ contains Γ_1. Then, the immersion $\mathbf{G}_1 \hookrightarrow \mathbf{G}$ induces an immersion $X_1 \hookrightarrow X$ whose image is a totally geodesic Riemann surface. (This follows, from example, from [**8**], Theorem 7.2, p. 224. See also [**12**], §4.1.)

Let us finally remark that in this example, M is a power of the Poincaré disk, hence its holomorphic sectional curvature is equal to $-1/\dim(M)$.

3.7. Hecke operators. — Any $g \in G(\mathbf{Q})$ gives rise to a *Hecke correspondence* T_g on X_Γ. Let us recall its definition. Let $\Gamma_g = \Gamma \cap g^{-1}\Gamma g$; this is a congruence subgroup of finite index in Γ. The variety $\Gamma_g\backslash M$ is smooth projective and admits two étale maps to X_Γ, given respectively by

$$p_1(\Gamma_g x) = \Gamma x, \qquad p_2(\Gamma_g x) = \Gamma g x,$$

for any $x \in M$, hence a correspondence on X_Γ.

Let us describe explicitly the image of a subvariety $Y \subset X_\Gamma$ by this correspondence. Let $(h_1, \dots, h_r)$ be a family of elements of Γ which represent each left-class modulo Γ_g. Let $\tilde{Y} \subset M$ be the preimage of Y; then,

$$T_g([Y]) = \sum_{i=1}^{r} [p(gh_i\tilde{Y})],$$

where $p \colon M \to X_G$ is the natural projection.

In view of this description, we observe that if Y is a totally geodesic Riemann surface in X_Γ. then so are all the irreducible components of $T_g([Y])$.

3.8. Weak approximation. — Since the group G is connected, $\mathbf{G}(\mathbf{Q})$ is dense in $G = \mathbf{G}(\mathbf{R})$ ([**13**], Theorem 7.7, p. 415). Let $\tilde{Y} \subset G$ be the preimage of Y by the projection map $G \to X_\Gamma = \Gamma\backslash G/K$. The union of the curves $a\tilde{Y}$, for $a \in G(\mathbf{Q})$, is therefore dense in G. It follows that the union of the curves $T_a Y$ is dense in X_Γ.

Let Y_0 be any fixed totally geodesic Riemann surface in X_Γ. We conclude that the irreducible components of the curves $T_a Y_0$, for $a \in G(\mathbf{Q})$, are dense in X_Γ. Since these components are also totally geodesic, we obtain the following result.

Theorem 8. — *Let X_Γ be a compact Shimura variety associated to a semisimple, connected and simply connected $\mathbf{Q}$-anisotropic algebraic group over $\mathbf{Q}$; let us assume that X_Γ contains a totally geodesic Riemann surface.*

Then, the set of algebraic curves Y in X_Γ such that

$$\deg_{K_{X_\Gamma}}(Y) = \left(\frac{-1}{S_{X_\Gamma}}\right)(-\chi(Y))$$

is dense in X_Γ for the complex topology; in particular, it is Zariski dense.

Remark 9. — When it applies, a theorem of Clozel, Oh and Ullmo [**6**] refines the density of Hecke orbits into an equidistribution theorem.

References

[1] D. Abramovich – "A linear lower bound on the gonality of modular curves", *Internat. Math. Res. Notices* (1996), no. 20, p. 1005–1011.

[2] A. L. Besse – *Einstein manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, 2008, Reprint of the 1987 edition.

[3] F. A. Bogomolov – "Families of curves on a surface of general type", *Dokl. Akad. Nauk SSSR* **236** (1977), no. 5, p. 1041–1044.

[4] J.-F. Boutot & H. Carayol – "Uniformisation p-adique des courbes de Shimura: les théorèmes de Čerednik et de Drinfel'd", Astérisque, no. 196-197, p. 7, 45–158 (1992), Astérisque **196-197**, Soc. Math. France, 1991.

[5] K. Buzzard – "Integral models of certain Shimura curves", *Duke Math. J.* **87** (1997), no. 3, p. 591–612.

[6] L. Clozel, H. Oh & E. Ullmo – "Hecke operators and equidistribution of Hecke points", *Invent. Math.* **144** (2001), no. 2, p. 327–351.

[7] J.-P. Demailly – "Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials", in *Algebraic geometry—Santa Cruz 1995*, Proc. Sympos. Pure Math., vol. 62, Amer. Math. Soc., Providence, RI, 1997, p. 285–360.

[8] S. Helgason – *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics, vol. 34, American Mathematical Society, Providence, RI, 2001, Corrected reprint of the 1978 original.

[9] M. McQuillan – "Non-commutative Mori theory", Preprint IHES/M/01/42, IHES, 2001.

[10] T. Miyake – *Modular forms*, english ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2006, Translated from the 1976 Japanese original by Yoshitaka Maeda.

[11] Y. Miyaoka – "The orbibundle Miyaoka-Yau-Sakai inequality and an effective Bogomolov-McQuillan theorem", *Publ. RIMS. Kyoto Univ.* **44** (2008), no. 2, p. 403–417.

[12] B. Moonen – "Linearity properties of Shimura varieties. I", *J. Algebraic Geometry* **7** (1998), no. 3, p. 539–567.

[13] V. Platonov & A. Rapinchuk – *Algebraic groups and number theory*, Pure and Applied Mathematics, vol. 139, Academic Press Inc., Boston, MA, 1994, Translated from the 1991 Russian original by Rachel Rowen.

[14] G. Shimura – *Introduction to the arithmetic theory of automorphic functions*, Publications of the Mathematical Society of Japan, Kanô Memorial Lectures, vol. 11, Princeton Univ. Press, 1994, Reprint of the 1971 original.

[15] Y. Tashiro – "On the holomorphic sectional curvature of the Bergman metric", *Sci. Rep. Tokyo Kyoiku Daigaku Sect. A* **8** (1963), p. 63–77 (1963).

[16] ———, "The curvature of the Bergman metric", *Sci. Rep. Tokyo Kyoiku Daigaku Sect. A* **8** (1965), p. 280–296 (1965).

[17] M.-F. Vignéras – *Arithmétique des algèbres de quaternions*, Lecture Notes in Math., no. 800, Springer-Verlag, 1980.

[18] P. Vojta – *Diophantine approximations and value distribution theory*, Lecture Notes in Mathematics, vol. 1239, Springer-Verlag, Berlin, 1987.

[19] ———, "A more general ABC conjecture", *Int. Math. Res. Not.* **21** (1998), p. 1103–1116.

Pascal Autissier, IMB, Université Bordeaux 1, 351 cours de la Libération, F-33405 Talence Cedex • *E-mail :* `pascal.autissier@math.u-bordeaux1.fr`

Antoine Chambert-Loir, Irmar, Université de Rennes 1, Campus de Beaulieu, F-35042 Rennes Cedex, France • *E-mail :* `antoine.chambert-loir@univ-rennes1.fr`

Carlo Gasbarri, IRMA, Université de Strasbourg, 7, Rue René Descartes, F-67000 Strasbourg • *E-mail :* `gasbarri@math.unistra.fr`